versus-host disease lethality in part via direct effects on CD4+ and CD8+ T cells," J. Immunol., 1996. Capon, et al., "Designing CD4 immunoadhesins for AIDS therapy," Nature, 1989, 337, 525—531 YR Cohen, J., New protein steals the show as 'costimulator' of T cells, Science, (1993) 262 844-845. ZR

PAGE 4/6 * RCVD AT 2/6/2004 3:22:55 PM [Eastern Standard Time] * SVR:USPTO-EFXRF-1/3 * DNS:8729306 * CSID:+703 905 2500 * DURATION (mm-ss):02-34

autoimmune encephalomyetitis," Eur. J. Immunol., 1981, 11, 195-189

PHULP GAMBEL 2/14/05

vR

WR

XR

Ben-Nun, et al. "The rapid isolation of clonable antigen-specific T lymphocyte lines capable of mediating

Blazar, et al., "Infusion of anti-B7.1 (CD80) and anti-B7.2 (CD86) monoclonal antibodies inhibits murine graft-

Feb-06-2004 15:24 From-PILLSBURY WINTHROP

+703-905-2500

T-868 P.005/006 F-500

	2004 15	:24 From-PILLSBURY WINTHKUP Trus-805-2000 1-000 P-000
10	AAR [Dautigny, et al., "Molecular cloning and nucleotide sequence of a cDNA clone coding for rat brain myelin proteolipid," FEBS Lett., 1985, 188(1):33-36.
4	BBR	Delabie, et al. The B7/BB1 antigen is expressed by Reed-Stetriberg colors." Blood, 1993, 82;2845-52.
T	DDR	Durie, et al., The role of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and its ligand (gp39) in peripheral and central control of CD40 and centra
1	EER	Durie, et al., "Prevention of collagen-induced armnus with an antibody to specific and agents."
	FFR	Falini, et al., "Response of refractory Hodgkin's disease to monocional and CD30 minhardes."
		Freeman, et al., "Uncovering of functional alternative CTLA-4 counter-receptor in D7-database with the D7-database with the counter-receptor in D7-database with the D7-database with the D7-database with the D7-database with the D7-database wi
	ннR	Freeman, et al., "Cloning of B7-2: a CTLA-4 counter-receptor that costimulates number 1 cell proliferation."
	IIR	Gernise, et al., "CD40-CD40 ligand interactions in experimental allergic enceptration years and managed
	אנג	Gottlieb, et al., "Results of a single-dose dose-escalating that of an anti-b/. T antibody (IDEC-1147 it) parasition of the control of the co
	KKR	Gottlieb, et al., "Clinical and histologic response to single-dose treatment of moderate in severe postations of moderate in severe postations." I Am. Acad. Dermatol. 2002, 47:693-700.
	LLR	Guinan, et al., "Pivotal role of the B7:CD28 pathway in transplantation tolerance and tumor transplantation
1	MMR	Hafter, et al., "The potential of restricted T cell recognition of myelin basis protein epitopes in the decapy of
	NNR	Hariharan, et al., 'Therapeutic activity of IDEC-114 (anti-CD80) and muximab (Killuxans) in 5-cell lymphonic
	OOR	Hathcock, et al., "Identification of an alternative CTLA-4 ligand costimulatory for 1 cell activation," Science,
1	PPR	Hollenbaugh, et al., "The human T cell antigen gp39, a member of the TNF gens family, is a lightly to the CD40 receptor expression of a soluble form of gp39 with B cell co-shmulatory activity," EMBO J., 1992, 11(12) 4313-4321
1	QQR	License Temporary Increase the Property of Curr Opin Immunol., 1992, 4:553-560.
1	RRR	Karpus, et a). "CD4+ suppressor cells differentially affect the production of in Ney by effective cells of
- -	629	The state has a strategies for drug design and discovery." Science, 1992, 257:1076-82.
_	SSR	Laman, et al., "The role of gp39 (CD40 ligand) in EAE and M5," Journal of Neuroinfinanciary, 1997, 547
T	UUR	1
	VVR	Lider, et al., "Suppression of experimental autoimmune encephalomyetius by oral administration of myelling et al., "Suppression of experimental autoimmune encephalomyetius by oral administration of myelling et al., "Suppression of experimental autoimmune encephalomyetius by oral administration of myelling et al.,"
	wwi	R Linsley, et al., "The role of the CD28 receptor during T cell responses to antigen," Artr. 1497 Interior, 13
	XXR	Liu. et al., "Blockade of CD28/CTLA-4-B7 co-stimulatory pathway in colfuc SCID mica. Digestive Disease
	YYR	McCafferty, et al., Phage antibodies. filamentous phage displaying antibody variable domains, Mature, 1
	ZZR	Miller, et al., "Antigen-driven bystander suppression after oral administration of antigens, "J Exp. Metc., 13
_	, ,	Mokhtarion, et al., "Adoptive transfer of myelin basic protein-sensitized 1 cells produces chiloric relapsing
		Morrison, et al., "Chimene human antibody molecules, mouse antigen-binding domains with human construction of Research Acad. Sci. USA, 1985, 81:6851-6855.
$\neg \uparrow$		Ifegion domains, Proc. read Acad. Sci. Sci. Sci. Sci. Sci. Sci. Sci. Sci
- 1		Tomos Stanturo Protection

PAGE 56 * RCVD AT 2/6/2004 3:22:55 PM [Eastern Standard Time] * SVR:USPTO-EFXRF-1/3 * DHIS:8729306 * CSID:+703 905 2500 * DURATION (mm-ss):02-34

PHILLIP GAMBEL 2/14/05

Feb-06-2004 15:24 From-PILLSBURY WINTHROP

+703-805-2500

T-868 P.006/006 F-500

	Z004 15	
4/1		Nickoloff, et al., "T tymphocytes in skin lesions of psoriasis and mycosis fungoides express B7-1; a ligand for CD28," <i>Blood</i> , 1994, 83:2580-2586.
-	FFFR	Noelle, et al., "A 39-kDa protein on activated helper T cells binds CD40 and transduces the signal for cognal
	GGGR	Oisson, et al., "Human-human monoclonal antibody-producing hybridomas, technical aspects," weth.
	НННВ	Pemn, et al., *Opposing effects of CTLA4-ig and anti-CD80 (B7-1) plus anti-CD86 (B7-2) on experimental effects of CTLA4-ig and anti-CD86 (B7-2) on experimental effects on the company of
1	IIIR	Pesoa, et al., Regulation of experimental allergic encephalomyelitis Part 5. Role of the recepient in
	אנונ	Pettinelli, et al. "Adoptive transfer of experimental allergic encephalomyelitis in SJD3 mice after in vision activation of lymph node calls by myelin basic protein: requirement for Lyt 1" 2" T lymphocytes," J. Immunol.
		Skolnick, et al., "From genes to protein structure and function: novel applications of computational structure and function in the structure and structure and function in the structure and
	1	Sobel, et al. "Acute experimental allergic encephalomyelitis in SJL/J mice induced by a synthetic peptide of
		Stamenkovic, et al., "A B-lymphocyte activation molecule related to the nerve growth ractor receptor and trade of the nerve growth ractor receptor receptor and trade of the nerve growth ractor receptor recept
	1	Suvas, et al Distinct role of CD80 and CD86 in the regulation of the activation of a cell and 5 cell
•	1	Takeda, et al., "Construction of chimaeric processed immunoglobulin genes containing mouse variable and process are process and process are process and process ar
		Teng, et al., "Construction and testing of mouse-human heteromyelomas for numan monocipital antibody
	l l	Tuony, et al., "Identification of an encephalitogenic determinant of myelin proteolipid protein for SUL mice,
		Van der Veen, et al., "The adoptive transfer of chronic relapsing experimental altergic encephalomyenus with more code cells sensitized to myelin proteoloid protein." <i>J. Neuroimmunol.</i> , 1989, 21:183-191.
	1	Ward, et al. "Binding activities of a repercire of single immunoglobulin variable domains secreted from
	В	Ward, et al., "Blocking of adhesion molecules in vivo as anti-inflammatory therapy," Their mundrion, 1999,
W	UUUF	1:105-171. R Yi-qun, et al. "Differential requirements for co-stimulatory signals from B7 family members by resting vers recently activated memory T cells towards soluble recall antigens," Int. Immunol., 1998, 8 37-44.
xamine		Date Considered:
		initial if citation considered, whether or not citation is in conformance with MPEP § 609. Draw line through conformance and not considered. Include copy of this form with next communication to Applicant.

PHURGANISA 1/14/05

FORM PI	ГО-144	9 (modified)	OTPE 3	Luc III		Atty. Ref. No 037003-0286			ent Ro 95-30	ef. -0233	CP3	
INFORMA BY APPL		DISCLOSURE S	•	A CONTRACTOR OF THE PARTY OF TH		Applicant:	ANDERSON	et al.				
012012						Appln. No.:	10/073,138					
						Filing Date:	February 13,	2002				
Date: Oct	ober 15	5, 2004	Page 1 of 2			Examiner:	Gambel, P.	Group Art	Unit:	. 164	44	
U.S. PAT	ENT D	OCUMENTS										
Examiner	s	Document	Date	Name				Class	Sub)	Filing	Date
Initials*		Number	MMYYYY	(Family N	ame of I	First Inventor	·)		Cla	ss	(if appro	opriate)
- pro-	AR	5,304,635	04/1994	lmam					+=		<u> </u>	-
	BR			ļ					1_		<u> </u>	
·	CR	<u> </u>						<u></u>	<u> </u>		<u> </u>	
FOREIGN	PATE	NT DOCUMENT	s · ·	••	* .	٠		English Abstract			slation ily Ava	
		Document	Date	Country	Inven	tor Name						
		Number	MM/YYYY		1			ļ		<u> </u>		-
	-							Enclosed	No	Enclo	se	No
	DR	ļ	_					 	<u> </u>			
	ER	 						 				
OT: 150 /	FR	<u> </u>				D. # 4.5		<u> </u>	-	<u> </u>		
		ig in this order Au						<u> </u>				<u></u>
M	GR	of a "Huma	nized" OKT3 i	monoclonal	antibod	y," J. Immun	the activating ar ol., 1992, 148:3	<u>461-3468.</u>				
	HR						Primary Alloge J. Exp. Med., 1				onses	and
	IR .	Azuma M., 177:845-85		onal Expres	sion of E	37/BB1 on A	ctivated T Lymp	hocytes," J	. Ехр	. Med	., 1993	3,
	JR		*	Antigen is a	a Secon	d Ligand for (CTLA-4 and CD	28," Nature	ə, 199	3, 36	3:76-7	9.
	KR	Boussiotis '	V.A., et al., "A	ctivated hun	man B ly	mphocytes e	express three C 1993, 90:11059	TLA-4 count 0-11063.	ter-re	ceptor	s that	co-
	LR						ne responses,"		men	t), Scie	ence, 1	1992,
	MR	De Boer M.	., et al., "Func y, 1992, 22:30		ecterizati	ion of a Nove	el Anti-B7 Mono	clonal Antib	ody,"	Eur.	Journa	al of
	NR				rsary for	r the war on o	cancer," Biotech	nology, 199	94, 12	2:320.		
	OR		, "The B7-2 (B 6 differentiatio				essed by mono 2-1407.	cytes and a	ctivte	d B lyr	nphoc	;ytes
	PR	Freeman G		LA-4 and C	D28 MF	RNA are Coe	xpressed in Mo	st T Cells A	fter A	ctivati	on," Th	he
	QR					_	nmune Tolerand	ce," Immuno	ology	Today	, 1993	3,
	RR	Gimmi C.D	o., et al., "Hum ion," Proc. Na				ed by Antigen F 90.	Presentation	in th	e Abs	ence c	of B7
	SR	Gribben J.					poptosis of hum	nan T cells."	Proc	. Nati.	Acad	. Sci.
	TŔ	Grumet F.0					Antigen Specific	ally Suppres	sses	Humo	ral	
	UR	Harding F.		28 Mediated	d Signal	ling Co-stimu	ılates Murine T	Cells and P	rever	nts Ind	uction	of

lonescu-Tirgoviste, et al, "Correlations between insulin antibodies and the HLA system in a group of Type I diabetic patients in Bucharest," Med. Interre, 1986, 24(1), 11-17. PHULP Grouper 2/14/05

Hart D.N.J., et al., "B7/BB-1 is a Leucocyte Differentiation Antigen on Human Dendritic Cells Induced by Activation." Immunology, 1993, 79:616-620

٧R

WR

<u>b</u>	XR -	Janeway C.A., Jr. and K. Bottomly, "Signals and Signs for Lymphocyte Responses," Cell, 1994, 76.275- 285.						
	YR	Jenkins M.K., "The Role of Cell Division in the Induction of Clonal Anergy." Immunology Today, 1992, 13:69.						
1	ZR	June C.H., et al., "The B7 and CD28 receptor families," Immunol Today, 1994, 15:321-331.						
	AAR	Krummel M., et al., "CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation," Exp. Med. 1995, 182:459-466.						
PE	BBR	LaSalle J.M., et al., "Early signaling defects in human T cells anergized by T cell presentation of autoantigen," J. Exp. Med., 1992, 176:177-186.						
97	ar Cil	Lenschow D.J., et al., "Long-Term Survival of Xenogeneic Pancreatic Islet Grafts Induced by CTLA-4Ig," Science, 1992, 257:789-795.						
THADS		Lenschow D.J., et al., "Expression and Functional Significance of an Additional Ligand for CTLA-4," ProcNatl. Acad. Sci., USA,,1993, 90:11054-11058.						
ТИНА В	EER	Lin H., et al., "Long-term Acceptance of Major Histocompatibility Complex Mismatched Cardiac Allografts Induced by CTLA-4-Ig Plus Donor Specific Transfusion," J. Exp. Med., 1993, 178:1801.						
-	FFR	Linsley, P.S., et al., "CTLA-4 is a Second Receptor for the B Cell Activation Antigen B7," J. Exp. Med., 1991, 174:561.						
	GGR	Linsley P.S., et. al., "T-Cell Antigen CD28 Mediates Adhesion with B Cells by Interacting with Activation Antigen B7/BB-1." Proc. Natl. Acad. Sci., 1990, 87:5031-5035.						
	HHR	Linsley, et al, "CD28 Engagement by B7/BB-1 Induces Transient Down-Regulation of CD28 Synthesis ar Prolonged Unresponsiveness to CD28 Signaling," The Journal of Immunology, 1993, 150:3161-3169.						
	IIR	Linsley, et al., "Binding of the B cell activation antigen B7 to CD28 costimulates T cell proliferation and interleukin 2 mRNA accumulation," J. Exp. Med., 1991, 173:721-730.						
	JJR	Linsley P.S., et. al., "Coexpression and Functional Cooperation of CTLA-4 and CD28 on Activated T Lymphocytes." J. Exp. Med., 1992, 176:1595-1604.						
	KKR	Morton P.A., et al., "Differential effects of CTLA-4 substitutions on the binding of human CD80 (B7-1) an CD86 (B7-2)," J. Immunol., 1996,156:1047-1054.						
	LLR	Nestle F.O., et al, "Characterization of dermal dendritic cells in psoriasis," J. Clin. Invest., 1994, 94: 202-209.						
	MMR	Schwartz R.H., "Co-stimulation of T lymphocytes: The role of CD28, CTLA-4, and B7/BB1 in interleukin-production and immunotherapy," Cell, 1992, 71:1065-1068.						
	NNR	Schwartz R.H., "A cell culture model for T lymphocyte clonal anergy," Science, 1990, 248:1349-1356.						
	OOR	Selvakumar,A., et al., "Genomic organization and chromosomal location of the human gene encoding the B-lymphocyte activation antigen B7," lmmunogenetics , 1992, 36:175-181						
	PPR	Tan P., et al., "Induction of Alloantigen-specific Hyporesponsiveness in Human T Lymphocytes by Blocki Interaction of CD28 with Its Natural Ligand B7/BB1," J. Exp. Med., 1993, 177:165-173.						
	QQR	Tivol E.A., et al., "Loss of CTLA-4 leads to massive lymphoproli-feration and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4," Immunity, 1995, 3:541-547.						
	RRR	Turka L.A., et al., "T-cell activation by the CD28 ligand B7 is required for cardiac allograft rejection in vivo						
	SSR	Toubert A., et al., "Epitope mapping of an HLA-B27 monoclonal antibody that also reacts with a 35-kD bacterial out-membrane protein," Clin. Exp. Immunol., 1990, 82(1), 16-20.						
	TTR	Toubert A., et al., "Epitope mapping of HLA-B27 and HLA-B7 antigens by using intradomain recombinan J. Immunol., 1988, 141(7), 2503-9.						
	UUR	Valle et al., "mAb 104, a new monoclonal antibody, recognizes the B7 antigen that is expressed on activated B cells and HTLV-1-transformed T cells," lmmunol. , 1990, 69(4), 531-535.						
	VVR	Vandenberghe P., et al "Antibody and B7/BB1-mediated ligation of the CD28 receptor induces tyrosine phosphorylation in human T cells," J. Exp. Med., 1992, 175:951-960.						
	WWR	van der Merwe, et al., "CD80 (B7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetic J. Exp. Med., 1997,185: 393-403.						
L	XXR	Weyl D., et al., "Epitope mapping of human monoclonal antibodies to HLA-B27 by using natural and mutated antigenic variants," Hum. Immunol., 1991, 31(4), 271-276.						
	YYR	Zavazava N., et al, "Inhibition of anti-HLA-B7 alloreactive CTL by affinity-purified soluble HLA,"						

*EXAMINER: Initial if citation considered, whether or not citation is in conformance with MPEP § 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to Applicant.

30486362_1.DOC